


AL12.2004-18

C.D.



## Fish & Wildlife Division

RESOURCE DATA AND  
SPECIES AT RISK SECTION

# Peace Parkland Native Grassland Stewardship Program 2001/02



Alberta Species at Risk Report No. 49





Digitized by the Internet Archive  
in 2016

[https://archive.org/details/peaceparklandnat00bake\\_0](https://archive.org/details/peaceparklandnat00bake_0)

# **Peace Parkland Native Grassland Stewardship Program 2001/02**

**Annette Baker**

**Alberta Species at Risk Report No. 49**

**2002**

**Project Partners**





Publication No.: I/070  
ISBN: 0-7785-2180-X (Printed Edition)  
ISBN: 0-7785-2181-8 (On-line Edition)  
ISSN: 1496-7219 (Printed Edition)  
ISSN: 1498-7146 (On-line Edition)

Cover Photograph: Native Grassland east of Grande Prairie by Annette Baker

For copies of this report, contact:

Information Centre – Publications  
Alberta Environment / Alberta Sustainable Resource Development  
Main Floor, Great West Life Building  
9920 108 Street  
Edmonton, Alberta, Canada T5K 2M4  
Telephone: (780) 422-2079

**OR**

Visit our web site at:

**<http://www3.gov.ab.ca/srd/fw/riskspecies/>**

This publication may be cited as:

Baker, A. 2002. Peace parkland native grassland stewardship program 2001/02. Alberta Sustainable Resource Development, Fish and Wildlife Division, Alberta Species at Risk Report No. 49. Edmonton, AB. 20 pp.

## DISCLAIMER

The views and opinions expressed are those of the author and do not necessarily represent the policies or positions of the Department or the Alberta Government.

## TABLE OF CONTENTS

|       |   |    |
|-------|---|----|
| 1.0   | INTRODUCTION .....                          | 1  |
| 1.1   | Goals and Objectives .....                  | 2  |
| 2.0   | STUDY AREA .....                            | 3  |
| 2.1   | General Description .....                   | 3  |
| 2.2   | Pilot Area for Method Comparison.....       | 3  |
| 2.3   | Vegetation.....                             | 4  |
| 3.0   | METHODS .....                               | 5  |
| 3.1   | Identifying Native Grasslands.....          | 5  |
| 3.2   | Field Inventory and Mapping.....            | 5  |
| 3.2.1 | Vegetation Inventory.....                   | 6  |
| 3.2.2 | Range Health.....                           | 6  |
| 3.2.3 | Butterflies .....                           | 6  |
| 3.2.4 | Rare Plants .....                           | 6  |
| 3.2.5 | Mapping.....                                | 7  |
| 3.3   | Stewardship.....                            | 7  |
| 4.0   | RESULTS .....                               | 8  |
| 4.1   | Location of Native Grassland Remnants ..... | 8  |
| 4.2   | Area of Native Grassland Remnants.....      | 8  |
| 4.3   | Vegetation Inventory .....                  | 8  |
| 4.4   | Range Health.....                           | 9  |
| 4.5   | Butterflies.....                            | 9  |
| 4.6   | Rare Plants .....                           | 9  |
| 4.7   | Stewardship.....                            | 9  |
| 4.8   | Mapping and Data Storage.....               | 10 |
| 5.0   | DISCUSSION AND FUTURE DIRECTIONS.....       | 11 |
| 6.0   | LITERATURE CITED .....                      | 13 |
|       | APPENDIX A.....                             | 15 |
|       | APPENDIX B.....                             | 20 |

## LIST OF FIGURES

|  |   |
|--|---|
| Figure 1. Northwest region of Alberta - Study and pilot areas of the Peace Native Grasslands Program, 2001. .... | 3 |
|--|---|

## LIST OF TABLES

|  |   |
|--|---|
| Table 1. Area (ha, %) and number (#, %) of upland native grassland remnants in each size category within the studied areas of Grande Prairie and Fairview, 2001 (n=43) ..... | 8 |
|--|---|



## ACKNOWLEDGEMENTS

To the landowners who gave access to their native grassland remnants, thank-you for the stewardship you already do, and for sharing these special sites with us. Darcy Meyers, Alberta Conservation Association, willingly assisted with fieldwork, data entry and anything else that was required, thank-you. Bonnie Hood, Resource Information Unit, Land and Forest Service, your professionalism, patience and attention to detail produced a quality of mapping that surpassed expectation. Anne Holcroft-Weerstra, Biota Consultants, without your expertise in air-photo interpretation this project would not have had a life beyond the first year.

This project was a combined effort of many individuals and organisations giving of their time, experience and enthusiasm in big, small and overlapping ways. Thanks to Ed Kolodychuk and Adam James for having the vision to start the program, and for helping out with subsequent project decisions, along with Carcey Rowand, Colin Stone, Margot Hervieux, Reg Arbuckle, Ron Bennett, and Tara Szkorupa. Bruce Sanders of Lands and Forest Service provided the use of air-photographs for interpretation. Brendan Kowalenko, Cameron Broatch, Chantel Winnicky, John Hallett, Mark Feldberg, Rene Labbe, Reg Arbuckle and Sheila Rangen helped with fieldwork. For botanical help, especially rare plants, thanks to Graham Griffiths, Joyce Gould, Ksenija Vujnovic, Lorna Allen, Patricia McIsaac and Sue Peters. Kim Banman and John Hallett checked data.

In 2001/02, funding was secured through the Government of Canada Habitat Stewardship Program for Species at Risk, the Alberta Government's Species at Risk program, and the Alberta Conservation Association. Support through in-kind contributions also came from Alberta Conservation Association, Alberta Fish and Wildlife Division, Alberta Lands and Forest Division, Alberta Public Lands Division, Ducks Unlimited Canada, and Peace Parkland Naturalists Club.

### Note:

Common plant names are from "Alberta Plants and fungi – Master species list and species group checklists" (Alberta Environmental Protection, 1993).

Butterfly names are from "Butterflies of Canada" (Layberry *et al*, 1998)



## EXECUTIVE SUMMARY

Historically, native grasslands were extensive and contiguous throughout the Peace Parkland of northwest Alberta. Land development has fragmented the landscape since 1900 and less than 0.5% of the original area of upland native grassland now remains. Land managers, researchers, writers and decision makers, often overlook these grasslands and there is little information on the location, area and state of remnants.

Native grasslands provide many benefits, including habitat for plant and animal species. In particular, a number of butterfly species with disjunct populations are found in the native grasslands of the Peace and they may be at risk from habitat disturbance and further habitat fragmentation.

A pilot project was initiated in 2001 to locate, inventory and map remaining native grasslands in two areas around Grande Prairie and Fairview. Air-photo interpretation provided the best means of locating small remnants and fieldwork included inventory of plant and butterfly species as well as assessment of range health. Remnants were mapped using Global Positioning Systems (GPS) and Geographical Information Systems (GIS) technology, which provided an electronic dataset, useful for planning.

As a result of the first year of this project, the documented area of native grasslands in the Peace River region has more than tripled. Conservation efforts should target the medium (10.1-30 ha) and large (30.1-65.0 ha) remnants first, as collectively they contribute the greatest area (87%), and should be easier to manage and maintain at a self-functioning size. The smaller remnants, which are the majority of sites, offer management challenges, which will require unique strategies to maintain them in their natural state.

In the future we will apply proven techniques of locating remnants from the pilot project to the entire white (settled) zone of the Northwest region. Continued work to further understand the relationship between butterflies and the native grassland on which they live, will be important to fill the current knowledge gaps that exist. The discovery of many new sites indicated that there is a need to work with landowners to conserve and maintain native grasslands that remain in the Peace Parkland region. Conservation of the sites through stewardship will become the primary means of protecting remnant native grasslands of the Peace Parkland region.

## 1.0 INTRODUCTION

Grasslands, which were once extensive and contiguous throughout the Peace River region, are now confined to small, scattered remnants in the uplands, and larger areas on south and west facing slopes of the Peace River and its tributaries (Alberta Environmental Protection 1997). Based on early survey data from Bezanson (1907), Kelly (1910), Macoun (1904), Mair (1908), and Wright (1930), cited in Wilkinson (1981), native grasslands covered an estimated 435,264 to 1,128,935 hectares in the Grande Prairie, Spirit River and Peace River areas. Less than a century later, the author estimates that less than 0.5% of the historical area of upland native grassland remains in the Peace River region.

Native grasslands were utilised for hay and pasture for cattle and horses as early as the late 1800s in the Spirit River area (Leonard 2000). The openness and rich soils under the grasslands ensured that these were the first areas to be cultivated when settlers arrived in the region in the early 1900s to homestead and establish widespread agriculture in the north (Leonard 2000). Towns and settlements scattered throughout the northwest testify to the importance of these open grasslands (prairies) through their names (e.g. Grande Prairie, High Prairie, Big Prairie, Salt Prairie, Paddle Prairie, John D'or Prairie, Buffalo Head Prairie, Wilson Prairie, Savage Prairie, Clear Prairie, Little Prairie, Prairie Echo, and Prairie Point).

The wide-scale loss of native grassland can be attributed to the expansion of cultivation and grazing, urbanisation, industrial development, encroachment of forests and shrub lands, and competition with non-native species (from tame pastures and invasive weeds). These factors have impacted all parkland areas of Alberta. However, the Peace River Parkland has undergone the most extensive alteration of all Alberta's parklands and fragmentation is recognised as the most serious threat to biological diversity within the region (Alberta Environmental Protection 1997).

Native grasslands are home to a diverse and interdependent mix of insects, birds, plants and mammals. The grasslands of the Peace River Parkland support a number of disjunct plant species (Alberta Environmental Protection 1997). Important relationships also exist between native grasslands and butterflies. For example, the larvae of the plains skipper (*Hesperia comma assiniboia*), Alberta arctic (*Oeneis alberta*) and Uhler's Arctic (*Oeneis uhleri varuna*) feed predominantly on native grasses and have disjunct populations in the Peace River area (Hervieux 2002). Native grasslands also provide habitat for vulnerable plant species like the leather grape fern (*Botrychium multifidum*; general status May be at Risk). Maintaining and conserving grasslands has the potential to conserve multiple species associated with or dependant upon these areas.

There has been little opportunity to study the grasslands of northern Alberta, and they are often overlooked in reference material and maps. The lack of information about native grasslands in the uplands of the Peace Parkland has also precluded their inclusion in recent descriptions and management recommendations of grasslands found in the Peace River Parkland, and these recommendations are restricted to the plant communities found



on the river slopes; e.g. "Guide to using native plants in Reclamation" (Gerling *et al.* 1999), and "Range Health Assessment" (Adams *et al.* 2000).

This project was initiated to increase knowledge about the Peace native grasslands. In particular, managers required information on the amount, location and condition of remnant patches, as well as the conservation threats and land use pressures. Using current Global Positioning System (GPS) and Geographical Information System (GIS) technology, conservation planning can be quickly visualised through the expediency and accuracy of spatially mapped information. This basic information will help to focus and coordinate the stewardship and conservation of native grasslands in the Peace region. In addition, data from the study of these remnants can be used by government, conservation organizations, reclamation industry and landowners for developing recommendations for management of these areas.

### 1.1 Goals and Objectives

The goal of the Peace Native Grassland Program was to identify and conserve native grassland remnants in the Northwest region. Specific objectives included, to:

1. Identify native grassland remnants by compiling existing information and identifying undocumented sites;
2. Inventory sites, collect vegetation information, and assess range health;
3. Document the presence of butterflies associated with native grassland habitats in association with the Peace Parkland Butterfly Survey.
4. Map remnants spatially, for use in GIS; and
5. Use information collected to direct stewardship activities:
  - a) Inform landowners of the conservation value of their property;
  - b) Provide information on the unique characteristics of native grasslands;
  - c) Provide information on the economic and social benefits of maintaining sites; and
  - d) Encourage landowners to participate in conservation easements, management agreements, land acquisition and other stewardship options to maintain areas.

## 2.0 STUDY AREA

### 2.1 General Description

The Northwest administrative region for Alberta Conservation Association (ACA) and Alberta Sustainable Resource Development covers 229,215 km sq, which is 34% of the provincial land base (Alberta Environment 2001). The region is divided into two land management zones: the White (settled) zone, primarily held under private ownership with some crown grazing leases, mineral leases and protected areas, and the Green (forestry) zone, primarily public land and, apart from some protected areas, is mainly used for logging, grazing and oil and gas operations. A variety of recreational activities also occur throughout the area.

This study considered all native grasslands throughout the region, but concentrated on the White zone as it was seen as being more at risk, had historical locations of grasslands, was held under private ownership and had existing resources such as air-photo coverage to enable a complete assessment. The White zone extends from Valleyview in the south to High Level in the north, and Slave Lake in the east to the Alberta/British Columbia border in the west. This area covers 43,203 km sq or 9% of the provincial land mass.

### 2.2 Pilot Area for Method Comparison

In 2001, the study concentrated on the Grande Prairie and Fairview areas. A pilot area within these areas was selected to determine the best method of locating remnants. Two areas, each approximately eight townships in size, were selected for the pilot. Each was known to contain native grasslands and was representative of the landscape found in the Fairview and Grande Prairie areas. Only the area in Grande Prairie was completed in 2001 due to the unavailability of air-photos for the Fairview area.




Figure 1. Northwest region of Alberta - Study and pilot areas of the Peace Native Grasslands Program, 2001.



### 2.3 Vegetation

Native grasslands contain a diverse mixture of grasses, forbs, and shrubs. Moss (1952) identified 36 species of grasses, sedges and rushes, 84 species of forbs and 19 species of woody plants for a total of 139 species of vascular plants in the Peace region. Native grasslands in the Peace are found on river slopes, uplands and lowlands and the plant assemblages differ under these various conditions. Our study focussed on upland native grasslands described by Moss (1952) as a wheat grass (*Agropyron*)- needle grass (*Stipa*) faciation of a wheat grass – needle grass – sedge (*Carex*) plant community. Other botanists have classified these native grasslands as sedge – intermediate oatgrass (*Danthonia intermedia*) – western porcupine grass (*Stipa spartea* var. *curtiseta*) communities (Wilkinson 1981) and (*Agropyron trachycaulum* var. *unilaterale*) – porcupine grass (*Stipa spartea*) – intermediate oat grass – sedge type (Wallis 1982). For a comprehensive overview of these community types see Vujnovic and Bentz (2001).

### 3.0 METHODS

#### 3.1 Identifying Native Grasslands

To locate remaining native grasslands, we used word of mouth, previous studies, government databases, happenstance, and air-photo interpretation. Air-photo interpretation gave the most number of potential sites and identified even small remnants. An experienced air-photo interpreter was contracted to interpret images for upland native grassland remnants in the White zone. The most recent available black and white air-photos, primarily at a scale of 1:30,000 (sometimes 1:40,000), were used to identify potential native grasslands. Interpretation was to 3-4 ha in size, or as small as could be discerned. Potential sites were outlined on a clear film 1:50,000 scale mapsheet that showed a 1998 Indian Resources Satellite (IRS) image and Alberta Township System (ATS) information. Completed mapsheets were used for field verification.

Historical descriptions from surveyors and pioneers were reviewed, and existing survey, land use, soil and vegetation maps were examined to see if this would identify targets on which to focus. While this gave a general idea of historical locations, the wide scale cultivation of the grassland since settlement has altered the landscape so much that these records could not accurately pinpoint the small remnants left today. In the Peace River region, solonchic soils play an important role in maintaining grasslands through the inhibition of tree root growth (Wilkinson 1981, Wilkinson and Johnson 1983). However soil conditions do not fully explain the location of all grasslands in the Peace (Wilkinson 1981, Wilkinson and Johnson 1983). The use of satellite imagery was also investigated, but as it is hard to reliably differentiate between tame and native grasses, this method was not recommended by similar studies being undertaken in Alberta's Grassland and Central Parkland natural regions (R. Bjorge *pers comm*).

#### 3.2 Field Inventory and Mapping

Often, researchers and rangeland specialists use indicator plant species or standard ecological site descriptions to determine plant community type. This information was very limited for upland native grasslands of the Peace Parkland and we first needed to gather the baseline data, which can be used to formulate these approaches. We also needed to verify the accuracy of the air-photo interpretation pilot. These needs were addressed through the collection of baseline data between June and August 2001.

We used the term "site" to refer to a quarter section (65 hectares in size). Within each site, "polygons" were open grassland areas separated by topography, vegetation or distance and each site could have multiple polygons. At each site, we familiarized ourselves with the polygons by traversing them, recording the presence of all plant species in the grassland and estimating plant species composition. If a polygon had more than 30% native species, it was classified as being native, if there was more than 70% non-native species it was classed as modified (Adams *et al.* 2000) and no further work was undertaken.



### 3.2.1 Vegetation Inventory

The most representative polygon of native grassland at each site was chosen for detailed vegetation sampling using the method described by Robertson and Adams (1990). The composition of plant species was estimated by listing each plant species present along with its cover estimate. Plants were identified to the species level where possible following the classification of Moss (1983). The cover estimate was determined using a Daubenmire cover class, which was assigned to each plant species, litter, exposed soil, and moss/lichen within each frame (plot). A 20 cm x 50 cm frame was used to delineate the sample area. Ten frames (or 5 if the site was extremely small) were sampled in the polygon. Frames were located at regular intervals measured along a traverse line, which generally cut through the middle of the polygon. Woody species growing in the grassland within the polygon were described using an estimated density distribution, height and percent area covered.

### 3.2.2 Range Health

To determine the ability of a site to perform certain functions, we assessed its health using the recently developed “Range/Pasture Health Assessment Short Form” (Adams *et al.* 2000). This assessment compares a site’s ecological processes (integrity and ecological status, community structure, hydrological function and nutrient cycling, site stability and noxious weeds) with a standard or reference plant community. The reference plant community is the potential natural community for the site under light grazing disturbance. There were limited standard plant communities to refer to for upland native grasslands of the Peace Parkland, so a basic description was formulated under the guidance of Colin Stone, Range Management Specialist, Public Lands, Peace River.

### 3.2.3 Butterflies

We recorded all butterflies seen, weather conditions and total survey time when visiting native grassland sites; these were the protocols developed by Hervieux (2001). We paid particular attention to observations of the plains skipper, Pike’s Old World swallowtail (*Papilio machaon pikei*), coral hairstreak (*Satyrrium titus*), striped hairstreak (*Satyrrium liparops*), Gorgone checkerspot (*Chlosyne gorgone carlota*), northern checkerspot (*Chlosyne palla*), Uhler’s arctic and Alberta arctic. We did not collect information when it was cloudy, extremely windy, or rainy so not all native grassland sites were inventoried for butterflies. Adult butterflies were identified by sight, using Bird *et al.* (1995), usually after capture with an insect net. Some individuals were collected for future DNA analysis by the University of Alberta and for a reference collection being developed by Alberta Fish and Wildlife. Data was submitted to the Peace Parkland Butterfly Survey and also sent to the Alberta Natural Heritage Information Centre (ANHIC).

### 3.2.4 Rare Plants

Prior to the field season, ANHIC provided all known occurrence records of tracked plants found in the Peace Parkland. Through consultation with ANHIC and Species at Risk

staff, we developed a list of rare and tracked plant species that might be found in this study. Under the current Alberta species risk assessments (Anon 2001), there was one “may-be-at-risk” species and 13 species whose status was “in-preparation” that were likely to be found in native grassland uplands of the Peace Parkland. During the traverse of the site and vegetation inventory, every effort was made to look in microhabitats that had the potential to support rare or tracked plant species. Observations of tracked plant species were recorded and submitted to ANHIC using their rare plant guidelines.

### 3.2.5 Mapping

A hand held GPS unit was used to record the location of each Daubenmire frame. The perimeter of the native grassland where the grassland met shrubs taller than 0.5 metres, or some other barrier (e.g. cultivation, fence-line, water body etc.) was traversed and recorded with the GPS. This gave an estimation of area and was incorporated into a GIS, which was used to map native grassland locations.

### 3.3 Stewardship

Landowner support was an important element of this project. After ownership of sites was determined, landowners were contacted for permission to enter their land and were invited to accompany staff during the field inspection. As follow-up to the visit, landowners were sent a copy of the list of plant species that were observed at the site.



## 4.0 RESULTS

### 4.1 Location of Native Grassland Remnants

Using the methods described above to locate potential remnants, 110 sites were identified within the Fairview and Grande Prairie areas. Of these, 83 (75%) sites were field checked and 43 (39%) sites were classified as having native upland grassland remnants. For various reasons, 4 sites were not accessed but an assessment “over the fence” was conducted and the results are included as part of the data set. The majority of native sites were concentrated around the Kleskun Hills, east of Grande Prairie. Three sites were located outside the method-comparison-pilot-area with one site located west of Grande Prairie near Saskatoon Lake and the other two south of Fairview. The location of these sites is considered sensitive information but will be made available to project partners for stewardship planning.

### 4.2 Area of Native Grassland Remnants

The native grassland remnants verified during 2001 covered a total area of 480 ha. The size of each remnant varied in size from 0.1 to 50.7 ha with an average size of 11.2 ha. Table 1 compares the area and number of sites in each size category. Nearly 60% (25) of the sites were less than 10.0 ha in size and while only 12% of the sites were in the 30.1-65.0 ha size range, these 5 sites represented almost half (45.8%) of the native grasslands. It also shows that the highest number of sites (38%) were under 3.0 ha but covered less than 4% (18.3 ha) of the total area.

Table 1. Area (ha, %) and number (#, %) of upland native grassland remnants in each size category within the studied areas of Grande Prairie and Fairview, 2001 ( $n=43$ )

| Remnant Size (ha)   | < 1.0      | 1.1 - 3.0   | 3.1 - 10.0  | 10.1 - 30.0   | 30.1 - 65.0   | Total |
|---------------------|------------|-------------|-------------|---------------|---------------|-------|
| Total area (ha)     | 1.9 (0.4%) | 16.4 (3.4%) | 43.0 (9.0%) | 198.8 (41.4%) | 219.6 (45.8%) | 479.7 |
| Number of Sites (#) | 8 (19%)    | 8 (19%)     | 9 (21%)     | 13 (30%)      | 5 (12%)       | 43    |

### 4.3 Vegetation Inventory

Detailed vegetation inventories were conducted on 37 (86%) of the 43 verified native upland grassland sites. The total number of species (shrubs, grasses, and forbs) that were identified through the inventory was 117. Of these, there were 69 native forbs, 26 native grasses and grass-like species, 15 native shrubs and 7 introduced forbs and grasses. An additional 61 vascular species were identified during the traverse portion of the site inspections.

Plant diversity at each native site ranged from 15 to 41 species with an average of 27. The occurrence of introduced grasses and forbs was generally quite low at each site (maximum of three and two respectively) and there were four sites with no introduced

species identified. Native forbs generally contributed the largest proportion of species with between 11 and 28 species and an average of 19 at each site. Native grasses in turn provided between 4 and 14 species with an average of 9 at each site. Up to 8 shrub species were identified at an individual site.

#### 4.4 Range Health

A range health assessment was done at each of the 37 inventoried native grassland sites. Sites could be rated as healthy, healthy with problems, or unhealthy. None of the sites were unhealthy, 32% (12 sites) were healthy with problems, and 68% (25 sites) were healthy. Approximately half (14 sites) of the healthy sites scored a perfect rating of 100%. Factors that affected the rating of sites included significant alterations in the plant community due to grazing or other disturbances, missing life form layers, and reduced amounts of litter. Most sites were stable with little evidence of accelerated erosion, and none of the sites had a noxious weed problem. Generally, the number of native forbs increased as range health improved and the number of grass species stayed the same, though the species and their abundance often changed. As a reference and comparison, range health was also assessed at eight other sites that were accessed (native shrub, slope, modified).

#### 4.5 Butterflies

Observations of 20 butterfly species were recorded at one third (13/37) of the inventoried upland native grassland sites, at one native shrub site and one modified grassland site. Two tracked species were observed; the plains skipper was recorded at five sites and the coral hairstreak at one site. The sites where these two species were observed had average or higher plant species diversity, a range health of healthy or healthy with problems, and were between 2.5 – 32.5 ha in area. Three other possibly disjunct species were also observed, the common ringlet (*Coenonympha inornata*), great spangled fritillary (*Speyeria cybele*) and garita skipper (*Oarisma garita*).

#### 4.6 Rare Plants

Leather grape fern, assessed as a “May be at Risk” fern in Alberta, was observed at eight native grassland sites and one modified grassland site. Most of these sites were grazed and, apart from the one modified grassland site, were in a healthy range condition (greater than 72% health score).

#### 4.7 Stewardship

Most of the sites were on privately owned land and we had the opportunity to speak with over 50 landowners. Only 4 of the 50 landowners withheld permission to access their land. Some landowners were sceptical about our interest in their native grassland. However, those we were able to speak with after completing the site assessment, expressed interest in the variety of plant species and in ways that they could maintain their sites.



#### 4.8 Mapping and Data Storage

An electronic and hardcopy map of potential and confirmed upland native grassland remnants has been produced and is available to project partners. The original is housed at ACA in Peace River. Information on the distribution of rare plants and uncommon butterflies is considered sensitive information and is not included here.

## 5.0 DISCUSSION AND FUTURE DIRECTIONS

When this project started, there were two commonly known and documented sites of remnant upland native grassland in the Grande Prairie area - Kleskun Hills Natural Area, and the Cochrane site. These two sites had a combined area of 143 ha. As a result of the first year of this project, the documented area of native grasslands in the Peace River region has more than tripled to 480 ha. More work is needed to complete the identification of remnants in the Grande Prairie and Fairview areas, and to expand the coverage to the rest of the White zone in the Peace River region will probably result in the identification of additional sites. To increase efficiency and maximize cost effectiveness of the air-photo interpretation, other lands of conservation interest (e.g., river breaks and wetlands) could be assessed at the same time.

Nearly all the native grassland remnants in the pilot area are concentrated within a ten-mile radius of the Kleskun Hills Natural Area, which makes this an important area to focus conservation and stewardship activities. While every site is important and valuable to conserve, efforts should target the medium (10.1-30 ha) and large (30.1-65.0 ha) remnants first, as they contribute the greatest area (87%), and should be easier to manage and maintain at a self-functioning size. The three largest sites are already protected as a Natural Area (Kleskun Hills Natural Area) or owned by a conservation organization (Cochrane Ranch). While other remnants may be protected in the future, they are currently under private ownership and are being utilized for agricultural production in the form of grazing or idle land. These sites offer a good opportunity to be maintained through voluntary landowner stewardship that may include management strategies like grazing or burning practices, and protection against land development, especially residential subdivision.

The majority of sites (59%) had remnants under 10.0 ha in size, which contributed to 13% of the total area. The medium-small (3.1-10.0 ha) remnants cover 9% of the area and due to an overall contribution and close proximity to larger remnants are important to maintain. The cost/area may be high due to the small area, but individual remnants would benefit from management strategies that can buffer them from encroachment of shrubs and introduced plants like awnless brome (*Bromus inermis*), and Kentucky bluegrass (*Poa pratensis*) while potentially increasing the area of native grassland.

Small remnants (1.1-2.0 ha) cover 3.4% of the area, and will hold the greatest challenge for conservation and management because they may be subject to the spread of introduced plants or native shrubs. The cost (financial and time) to maintain these small sites could be prohibitive. However, as long as they are intact, they contribute to local biodiversity and may provide important habitat for some species of butterflies (M. Hervieux, *pers comm.*).

The smallest sites (<1.0 ha) only cover 0.4% of the area and if they are not close to one another to provide effective habitat for butterflies, may not be cost or time effective to secure. However, even these small sites may be important as sources of genetic material to help in the conservation of native grasslands in the Peace River region.



Landowners were interested in the results of this study, especially information concerning plant species that occurred on their property. Landowners also had questions about ways to best manage the native grassland, and why they are so important. A brochure or fact sheet that provides this information would help landowners and promote conservation of native grasslands. A short questionnaire about the landowners' interest in conservation is planned for 2002/2003 to help facilitate land stewardship efforts.

Vegetation data collected through the inventory has been helpful to gain a broader picture of the plant composition of the upland remnants. Further inventories at more sites, will increase this knowledge and can be used to make more accurate descriptions of reference plant community types. It will also give a better sense of the overall diversity of native grassland plant communities growing in the Peace River region.

There is limited knowledge about the food and nectar sources required for individual butterfly species. Further work on plant community types that support particular butterfly species may bridge this gap, and potentially, indicator species could be used to predict butterfly occurrences. This would be especially helpful as butterflies vary within a season (flight time and climatic conditions) and between seasons, so it is difficult to observe all species on a single visit. Documenting the phenology of plants and corresponding butterflies present may also help with this analysis and will be incorporated in future years.

The discovery of many overlooked remnants indicates that there is a need to work with landowners to conserve these areas and to inform people that native grasslands are an important part of the Peace River region landscape.

## 6.0 LITERATURE CITED

- Adams, B.W., A. Robertson, M. Willoughby, G. Ehlert, M. Alexander, D. Downing, D. Lawrence, C. Lane, and C. Stone. October 17, 2000. Range/pasture health assessment short form. Preliminary report for the Alberta Rangeland Health Task Group, Alberta, Canada.
- Alberta Environment. May 2001. URL:  
<http://www.gov.ab.ca/env/regions/nwb/index.html>
- Alberta Environmental Protection. 1993. Alberta Plants and fungi – Master species list and species group checklists. Edmonton, Alberta, Canada.
- Alberta Environmental Protection. 1997. The parkland natural region of Alberta: One of a series of reports prepared for the Special Places 2000 Provincial Coordinating Committee.
- Anonymous. 2001. The General Status of Alberta Wild Species 2000. Alberta Fish and Wildlife, Sustainable Resource Development, Edmonton. URL:  
<http://www3.gov.ab.ca/srd/fw/status/index.html>
- Bird, C.D., G.J. Hilchie, N.G. Kondla, E.M. Pike and F.A.H. Sperling. 1995. Alberta butterflies. The Provincial Museum of Alberta, Edmonton, Alberta, Canada.
- Gerling, H.S., M.G. Willoughby, A. Schoepf, E.E. Tannas and C.A. Tannas. 1996. A guide to using native plants on disturbed lands. Alberta Agriculture, Food and Rural Development and Alberta Environmental Protection.
- Hervieux, M. 2002. Survey of native grassland butterflies in the Peace parkland region of northwestern Alberta – 2001. Alberta Sustainable Resource Development, Fish and Wildlife Division, Alberta Species at Risk Report No. 47. Edmonton, Alberta, Canada.
- Layberry, R.A., P.W. Hall and J.D. Lafontaine. 1998. The butterflies of Canada. University of Toronto Press, Toronto, Canada.
- Leonard, D.W. 2000. Delayed frontier – The Peace Country to 1909. Detselig Enterprises Ltd., Calgary, Alberta, Canada
- Moss, E.H. 1952. Grassland of the Peace River region, western Canada. Canadian Journal of Botany 30: 98-124.
- Moss, E.H. 1983. Flora of Alberta, 2<sup>nd</sup> edition. University of Toronto Press Inc., Toronto, Ontario, Canada.



- Robertson, A., and B.W. Adams. 1990. Two worksheets for range vegetation monitoring. Range Notes Issue No. 8. Alberta Forestry, Lands and Wildlife, Public Lands Division, Alberta, Canada.
- Vujnovic, K., and J. Bentz. (Geowest Environmental Consultants Ltd.) 2001. Preliminary classification of native wheat grass (*Agropyron* spp.) community types in Alberta. Report prepared for Alberta Natural Heritage Information Centre, Alberta Environment, Edmonton, Alberta, Canada
- Wallis, C.A. 1982. An overview of the mixed grasslands of North America. Grassland Ecology and Classification: Symposium proceeding June 1982. Ministry of Forests, British Columbia, Canada.
- Wilkinson, K. 1981. Remnant and early settlement prairies and solonetzic soils in the Peace River district. MSc Thesis, University of Calgary, Calgary, Alberta, Canada.
- Wilkinson, K., and E.A. Johnson. 1983. Distribution of prairies and solonetzic soils in the Peace River district, Alberta. Canadian Journal of Botany 61: 1851-1860

## APPENDIX A.

### Plants identified on upland native grasslands remnants, in the Peace region of Northwest Alberta, by the *Peace Native Grassland Program*, 2001.

| <u>Scientific Name</u>                  | <u>Common name</u>         | <u>Family</u>    |
|---|----------------------------|------------------|
| <b>FORBS</b>                            |                            |                  |
| <i>Achillea millefolium</i>             | common yarrow              | Compositae       |
| * <i>Agoseris glauca</i>                | yellow false dandelion     | Compositae       |
| * <i>Agrimonia striata</i>              | agrimony                   | Rosaceae         |
| <i>Allium cernuum</i>                   | nodding onion              | Liliaceae        |
| * <i>Allium schoenoprasum</i>           | wild chives                | Liliaceae        |
| <i>Allium textile</i>                   | prairie onion              | Liliaceae        |
| <i>Androsace septentrionalis</i>        | northern fairy candelabra  | Primulaceae      |
| <i>Anemone</i>                          | anemone                    | Ranunculaceae    |
| <i>Anemone canadensis</i>               | Canada anemone             | Ranunculaceae    |
| <i>Anemone cylindrica</i>               | long-fruited anemone       | Ranunculaceae    |
| * <i>Anemone multifida</i>              | cut-leaved anemone         | Ranunculaceae    |
| <i>Anemone patens</i>                   | prairie crocus             | Ranunculaceae    |
| <i>Antennaria</i>                       | Everlasting                | Compositae       |
| <i>Antennaria neglecta</i>              | broad-leaved everlasting   | Compositae       |
| <i>Antennaria parvifolia</i>            | small-leaved everlasting   | Compositae       |
| * <i>Arabis</i>                         | rock cress                 | Cruciferae       |
| <i>Arnica</i>                           | arnica                     | Compositae       |
| * <i>Arnica chamissonis</i>             | leafy arnica               | Compositae       |
| <i>Arnica cordifolia</i>                | heart-leaved arnica        | Compositae       |
| <i>Arnica fulgens</i>                   | shining arnica             | Compositae       |
| <i>Artemisia</i>                        | artemisia                  | Compositae       |
| <i>Artemisia biennis</i>                | biennial sagewort          | Compositae       |
| <i>Artemisia campestris</i>             | plains wormwood            | Compositae       |
| <i>Artemisia frigida</i>                | pasture sagewort           | Compositae       |
| <i>Artemisia ludoviciana</i>            | prairie sagewort           | Compositae       |
| <i>Aster</i>                            | aster                      | Compositae       |
| <i>Aster conspicuus</i>                 | showy aster                | Compositae       |
| <i>Aster ericoides</i>                  | tufted white prairie aster | Compositae       |
| <i>Aster laevis</i>                     | smooth aster               | Compositae       |
| <i>Astragalus</i>                       | milk vetch                 | Leguminosae      |
| * <i>Atriplex nuttallii</i> (from Moss) | Nuttall's atriplex         | Chenopodiaceae   |
| <i>Botrychium multifidum</i>            | leather grape fern         | Ophioglossaceae  |
| * <i>Brassica</i>                       | mustard                    | Cruciferae       |
| <i>Campanula rotundifolia</i>           | harebell                   | Campanulaceae    |
| * <i>Capsella bursa-pastoris</i>        | shepherd's-purse           | Cruciferae       |
| <i>Castilleja miniata</i>               | common red paintbrush      | Scrophulariaceae |
| <i>Cerastium</i>                        | chickweed                  | Caryophyllaceae  |
| <i>Cerastium arvense</i>                | field mouse-ear chickweed  | Caryophyllaceae  |
| * <i>Chenopodium album</i>              | lamb's-quarters            | Chenopodiaceae   |

|                                    |                            |                  |
|------------------------------------|----------------------------|------------------|
| * <i>Cicuta maculata</i>           | water-hemlock              | Umbelliferae     |
| * <i>Cirsium</i>                   | thistle                    | Compositae       |
| <i>Cirsium drummondii</i>          | Drummond's thistle         | Compositae       |
| <i>Comandra umbellata</i>          | bastard toadflax           | Santalaceae      |
| * <i>Cornus canadensis</i>         | bunchberry                 | Cornaceae        |
| * <i>Delphinium glaucum</i>        | tall larkspur              | Ranunculaceae    |
| * <i>Descurainia sophia</i>        | flixweed                   | Cruciferae       |
| * <i>Dracocephalum parviflorum</i> | American dragonhead        | Labiatae         |
| * <i>Epilobium angustifolium</i>   | common fireweed            | Onagraceae       |
| * <i>Equisetum</i>                 | horsetail                  | Equisetaceae     |
| <i>Erigeron</i>                    | fleabane                   | Compositae       |
| <i>Erigeron caespitosus</i>        | tufted fleabane            | Compositae       |
| <i>Erigeron glabellus</i>          | smooth fleabane            | Compositae       |
| <i>Erigeron philadelphicus</i>     | Philadelphia fleabane      | Compositae       |
| <i>Fragaria virginiana</i>         | wild strawberry            | Rosaceae         |
| <i>Galium boreale</i>              | northern bedstraw          | Rubiaceae        |
| * <i>Gentianella</i>               | gentian/felwort            | Gentianaceae     |
| <i>Gentianella amarella</i>        | felwort                    | Gentianaceae     |
| <i>Geum triflorum</i>              | three-flowered avens       | Rosaceae         |
| <i>Grindelia squarrosa</i>         | gumweed                    | Compositae       |
| * <i>Habenaria</i>                 | bog orchid                 | Orchidaceae      |
| <i>Hedysarum alpinum</i>           | alpine hedysarum           | Leguminosae      |
| * <i>Helianthus</i>                | sunflower                  | Compositae       |
| * <i>Helianthus annuus</i>         | common annual sunflower    | Compositae       |
| * <i>Heracleum lanatum</i>         | cow parsnip                | Umbelliferae     |
| <i>Heuchera richardsonii</i>       | Richardson's alumroot      | Saxifragaceae    |
| <i>Hieracium umbellatum</i>        | narrow-leaved hawkweed     | Compositae       |
| <i>Lathyrus ochroleucus</i>        | cream-colored vetchling    | Leguminosae      |
| <i>Lilium philadelphicum</i>       | western wood lily          | Liliaceae        |
| <i>Linum lewisii</i>               | wild blue flax             | Linaceae         |
| <i>Maianthemum canadense</i>       | wild lily-of-the-valley    | Liliaceae        |
| * <i>Matricaria perforata</i>      | scentless chamomile        | Compositae       |
| * <i>Medicago sativa</i>           | alfalfa                    | Leguminosae      |
| * <i>Melilotus officinalis</i>     | yellow sweet-clover        | Leguminosae      |
| * <i>Mertensia paniculata</i>      | tall lungwort              | Boraginaceae     |
| <i>Monarda fistulosa</i>           | wild bergamot              | Labiatae         |
| * <i>Opuntia fragilis</i>          | brittle prickly-pear       | Cactaceae        |
| <i>Orthocarpus luteus</i>          | owl-clover                 | Scrophulariaceae |
| <i>Oxytropis</i>                   | locoweed                   | Leguminosae      |
| * <i>Oxytropis sericea</i>         | early yellow locoweed      | Leguminosae      |
| <i>Oxytropis splendens</i>         | showy locoweed             | Leguminosae      |
| * <i>Penstemon gracilis</i>        | lilac-flowered beardtongue | Scrophulariaceae |
| <i>Penstemon procerus</i>          | slender blue beardtongue   | Scrophulariaceae |
| * <i>Plantago major</i>            | common plantain            | Plantaginaceae   |
| <i>Potentilla</i>                  | cinquefoil                 | Rosaceae         |
| <i>Potentilla arguta</i>           | white cinquefoil           | Rosaceae         |



|                                   |                              |                 |
|-----------------------------------|------------------------------|-----------------|
| <i>Potentilla gracilis</i>        | graceful cinquefoil          | Rosaceae        |
| <i>Potentilla hippiana</i>        | woolly cinquefoil            | Rosaceae        |
| * <i>Ranunculus</i>               | buttercup                    | Ranunculaceae   |
| * <i>Ranunculus acris</i>         | tall buttercup               | Ranunculaceae   |
| <i>Ranunculus macounii</i>        | Macoun's buttercup           | Ranunculaceae   |
| * <i>Rumex</i>                    | dock                         | Polygonaceae    |
| <i>Sedum lanceolatum</i>          | lance-leaved stonecrop       | Crassulaceae    |
| <i>Selaginella densa</i>          | prairie selaginella          | Selaginellaceae |
| * <i>Senecio eremophilus</i>      | cut-leaved ragwort           | Compositae      |
| <i>Sisyrinchium montanum</i>      | common blue-eyed grass       | Iridaceae       |
| <i>Smilacina stellata</i>         | star-flowered Solomon's seal | Liliaceae       |
| <i>Solidago</i>                   | goldenrod                    | Compositae      |
| * <i>Solidago canadensis</i>      | Canada goldenrod             | Compositae      |
| <i>Solidago missouriensis</i>     | low goldenrod                | Compositae      |
| <i>Solidago spathulata</i>        | mountain goldenrod           | Compositae      |
| * <i>Sonchus</i>                  | sow thistle                  | Compositae      |
| * <i>Spiranthes romanzoffiana</i> | hooded ladies'-tresses       | Orchidaceae     |
| <i>Stellaria</i>                  | chickweed                    | Caryophyllaceae |
| * <i>Tanacetum vulgare</i>        | common tansy                 | Compositae      |
| <i>Taraxacum officinale</i>       | common dandelion             | Compositae      |
| <i>Thalictrum venulosum</i>       | veiny meadow rue             | Ranunculaceae   |
| * <i>Thlaspi arvense</i>          | stinkweed                    | Cruciferae      |
| * <i>Tragopogon dubius</i>        | common goat's-beard          | Compositae      |
| <i>Trifolium</i>                  | clover                       | Leguminosae     |
| <i>Trifolium hybridum</i>         | alsike clover                | Leguminosae     |
| * <i>Trifolium pratense</i>       | red clover                   | Leguminosae     |
| * <i>Trifolium repens</i>         | white clover                 | Leguminosae     |
| * <i>Urtica dioica</i>            | common nettle                | Urticaceae      |
| <i>Vicia americana</i>            | wild vetch                   | Leguminosae     |
| * <i>Viola canadensis</i>         | western Canada violet        | Violaceae       |
| <i>Viola adunca</i>               | early blue violet            | Violaceae       |
| * <i>Zizia aptera</i>             | heart-leaved Alexanders      | Umbelliferae    |

## GRASSES and GRASS-LIKES

|  |                        |           |
|--|------------------------|-----------|
| <i>Agropyron</i>   | wheat grass            | Gramineae |
| <i>Agropyron dasystachyum</i>                            | northern wheat grass   | Gramineae |
| <i>Agropyron repens</i>                                  | quack grass            | Gramineae |
| <i>Agropyron smithii</i>                                 | western wheat grass    | Gramineae |
| <i>Agropyron trachycaulum</i>                            | slender wheat grass    | Gramineae |
| <i>Agropyron trachycaulum</i> var.<br><i>unilaterale</i> | awned wheat grass      | Gramineae |
| <i>Agrostis scabra</i>                                   | rough hair grass       | Gramineae |
| * <i>Avena fatua</i>                                     | wild oat               | Gramineae |
| * <i>Bromus ciliatus</i>                                 | fringed brome          | Gramineae |
| <i>Bromus inermis</i>                                    | awnless brome          | Gramineae |
| * <i>Bromus pumpellianus</i> (from Moss)                 | northern awnless brome | Gramineae |

|                                   |                         |            |
|-----------------------------------|-------------------------|------------|
| <i>Calamagrostis</i>              | reed grass              | Gramineae  |
| * <i>Calamagrostis canadensis</i> | bluejoint               | Gramineae  |
| <i>Calamagrostis inexpansa</i>    | northern reed grass     | Gramineae  |
| <i>Calamagrostis montanensis</i>  | plains reed grass       | Gramineae  |
| <i>Calamagrostis purpurascens</i> | purple reed grass       | Gramineae  |
| <i>Carex</i>                      | sedge                   | Cyperaceae |
| <i>Danthonia intermedia</i>       | intermediate oat grass  | Gramineae  |
| <i>Deschampsia cespitosa</i>      | tufted hair grass       | Gramineae  |
| * <i>Elymus innovatus</i>         | hairy wild rye          | Gramineae  |
| * <i>Festuca rubra</i>            | red fescue              | Gramineae  |
| <i>Festuca saximontana</i>        | Rocky Mountain fescue   | Gramineae  |
| <i>Helictotrichon hookeri</i>     | Hooker's oat grass      | Gramineae  |
| * <i>Hierochloa odorata</i>       | sweet grass             | Gramineae  |
| * <i>Hordeum jubatum</i>          | foxtail barley          | Gramineae  |
| <i>Juncus</i>                     | rush                    | Juncaceae  |
| <i>Koeleria macrantha</i>         | June grass              | Gramineae  |
| <i>Muhlenbergia richardsonis</i>  | mat muhly               | Gramineae  |
| <i>Phleum pratense</i>            | timothy                 | Gramineae  |
| <i>Poa</i>                        | bluegrass               | Gramineae  |
| <i>Poa compressa</i>              | Canada bluegrass        | Gramineae  |
| * <i>Poa interior</i>             | inland bluegrass        | Gramineae  |
| <i>Poa palustris</i>              | fowl bluegrass          | Gramineae  |
| <i>Poa pratensis</i>              | Kentucky bluegrass      | Gramineae  |
| <i>Poa sandbergii</i>             | Sandberg bluegrass      | Gramineae  |
| <i>Schizachne purpurascens</i>    | purple oat grass        | Gramineae  |
| * <i>Scirpus</i>                  | bulrush                 | Cyperaceae |
| * <i>Stipa columbiana</i>         | Columbia needle grass   | Gramineae  |
| * <i>Stipa comata</i>             | needle-and-thread       | Gramineae  |
| <i>Stipa curtisetia</i>           | western porcupine grass | Gramineae  |
| <i>Stipa richardsonii</i>         | Richardson needle grass | Gramineae  |
| <i>Stipa viridula</i>             | green needle grass      | Gramineae  |
| * <i>Triticum aestivum</i>        | common wheat            | Gramineae  |
| * <i>Typha latifolia</i>          | common cattail          | Typhaceae  |

## SHRUBS

|                                |                     |                |
|--------------------------------|---------------------|----------------|
| * <i>Alnus</i>                 | alder               | Betulaceae     |
| <i>Amelanchier alnifolia</i>   | saskatoon           | Rosaceae       |
| <i>Arctostaphylos uva-ursi</i> | common bearberry    | Ericaceae      |
| <i>Cornus stolonifera</i>      | red-osier dogwood   | Cornaceae      |
| <i>Elaeagnus commutata</i>     | silverberry         | Elaeagnaceae   |
| <i>Juniper</i>                 | Juniper             | Cupressaceae   |
| * <i>Juniperus communis</i>    | ground juniper      | Cupressaceae   |
| <i>Juniperus horizontalis</i>  | creeping juniper    | Cupressaceae   |
| * <i>Lonicera dioica</i>       | twining honeysuckle | Caprifoliaceae |
| <i>Lonicera involucrata</i>    | bracted honeysuckle | Caprifoliaceae |
| <i>Populus tremuloides</i>     | aspen               | Salicaceae     |

|                                    |                     |                 |
|------------------------------------|---------------------|-----------------|
| <i>Ribes oxycanthoides</i>         | northern gooseberry | Grossulariaceae |
| <i>Ribes triste</i>                | wild red currant    | Grossulariaceae |
| <i>Rosa</i>                        | rose                | Rosaceae        |
| <i>Rubus idaeus</i>                | wild red raspberry  | Rosaceae        |
| <i>Salix</i>                       | willow              | Salicaceae      |
| <i>Shepherdia canadensis</i>       | Canada buffaloberry | Elaeagnaceae    |
| <i>Symphoricarpos occidentalis</i> | buckbrush           | Caprifoliaceae  |
| * <i>Vaccinium myrtilloides</i>    | common blueberry    | Ericaceae       |

\* identified during the traverse, but were not recorded in the vegetation inventory

Source used for common names:

Alberta Environmental Protection. 1993. *Alberta Plants and Fungi - Master Species List and Species Group Checklists*. Edmonton.



## APPENDIX B.

### Butterflies observed by *Peace Native Grasslands Program*, in the Peace River region during 2001

#### Skippers

|                   |                                  |
|-------------------|----------------------------------|
| *Plains Skipper   | <i>Hesperia comma assiniboia</i> |
| **Garita Skipper  | <i>Oarisma garita</i>            |
| Long-dash Skipper | <i>Polites mystic</i>            |

#### Swallowtails

Swallowtail sp.

#### Whites, Marbles and Sulphurs

|               |                     |
|---------------|---------------------|
| Cabbage White | <i>Pieris rapae</i> |
| White sp.     |                     |
| Sulphur sp.   |                     |

#### Coppers, Hairstreaks and Blues

|                   |                        |
|-------------------|------------------------|
| *Coral Hairstreak | <i>Satyrrium titus</i> |
| Northern Blue     | <i>Lycaeides idas</i>  |
| Blue sp.          |                        |

#### Tortoiseshells, Admirals and Anglewings

|               |                           |
|---------------|---------------------------|
| White Admiral | <i>Limenitis arthemis</i> |
|---------------|---------------------------|

#### Fritillaries

|                             |  |
|-----------------------------|--|
| Meadow Fritillary           | <i>Boloria bellona</i>                   |
| Aphrodite Fritillary        | <i>Speyeria aphrodite</i>                |
| Northwestern Fritillary     | <i>Speyeria electa</i>                   |
| **Great Spangled Fritillary | <i>Speyeria cybele pseudocarpenterii</i> |
| Mormon Fritillary           | <i>Speyeria mormonia</i>                 |
| Greater Fritillary sp.      |  |

#### Checkerspots and Crescents

|                         |                         |
|-------------------------|-------------------------|
| Northern Pearl Crescent | <i>Phyciodes cocyta</i> |
|-------------------------|-------------------------|

#### Satyrns

|                             |                             |
|-----------------------------|-----------------------------|
| Common Wood Nymph           | <i>Cercyonis pegala</i>     |
| **Common (Inornate) Ringlet | <i>Coenonympha inornata</i> |

\* = tracked butterflies found in Peace River region

\*\* = disjunct Alberta butterflies found in Peace River region

## 7.0 LIST OF TITLES IN THIS SERIES

(as of February 2004)

- No. 1 Alberta species at risk program and projects 2000-2001, by Alberta Sustainable Resource Development, Fish and Wildlife Division. (2001)
- No. 2 Survey of the peregrine falcon (*Falco peregrinus anatum*) in Alberta, by R. Corrigan. (2001)
- No. 3 Distribution and relative abundance of the shortjaw cisco (*Coregonus zenithicus*) in Alberta, by M. Steinhilber and L. Rhude. (2001)
- No. 4 Survey of the bats of central and northwestern Alberta, by M.J. Vonhof and D. Hobson. (2001)
- No. 5 2000 survey of the Trumpeter Swan (*Cygnus buccinator*) in Alberta, by M.L. James and A. James. (2001)
- No. 6 2000/2001 Brassy Minnow inventory at Musreau Lake and outlet, by T. Ripley. (2001)
- No. 7 Colonial nesting waterbird survey in the Northwest Boreal Region – 2000, by M. Hanneman and M. Heckbert. (2001)
- No. 8 Burrowing owl trend block survey and monitoring - Brooks and Hanna areas, by D. Scobie and R. Russell. (2000)
- No. 9 Survey of the Lake Sturgeon (*Acipenser fulvescens*) fishery on the South Saskatchewan River, Alberta (June-September, 2000), by L.A. Winkel. (2000)
- No. 10 An evaluation of grizzly bear-human conflict in the Northwest Boreal Region of Alberta (1991-2000) and potential mitigation, by T. Augustyn. (2001)
- No. 11 Harlequin duck monitoring in the Northern East Slopes of Alberta: 1998-2000 preliminary results, by J. Kneteman and A. Hubbs. (2000)
- No. 12 Distribution of selected small mammals in Alberta, by L. Engley and M. Norton. (2001)
- No. 13 Northern leopard frog reintroduction. Raven River - Year 2 (2000), by K. Kendell. (2001)
- No. 14 Cumulative effects of watershed disturbances on fish communities in the Kakwa and Simonette watersheds. The Northern Watershed Project. Study 3 Progress report, by T. Thera and A. Wildeman. (2001)
- No. 15 Harlequin duck research in Kananaskis Country in 2000, by C.M. Smith. (2001)
- No. 16 Proposed monitoring plan for harlequin ducks in the Bow Region of Alberta, by C.M. Smith. (2001)

- No. 17 Distribution and relative abundance of small mammals of the western plains of Alberta as determined from great horned owl pellets, by D. Schowalter. (2001)
- No. 18 Western blue flag (*Iris missouriensis*) in Alberta: a census of naturally occurring populations for 2000, by R. Ernst. (2000)
- No. 19 Assessing chick survival of sage grouse in Canada, by C.L. Aldridge. (2000)
- No. 20 Harlequin duck surveys of the Oldman River Basin in 2000, by D. Paton. (2000)
- No. 21 Proposed protocols for inventories of rare plants of the Grassland Natural Region, by C. Wallis. (2001)
- No. 22 Utilization of airphoto interpretation to locate prairie rattlesnake (*Crotalus viridis viridis*) hibernacula in the South Saskatchewan River valley, by J. Nicholson and S. Rose. (2001)
- No. 23 2000/2001 Progress report on caribou research in west central Alberta, by T. Szkorupa. (2001)
- No. 24 Census of swift fox (*Vulpes velox*) in Canada and Northern Montana: 2000-2001, by A. Moehrensclager and C. Moehrensclager. (2001)
- No. 25 Population estimate and habitat associations of the long-billed curlew in Alberta, by E.J. Saunders. (2001)
- No. 26 Aerial reconnaissance for piping plover habitat in east-central Alberta, May 2001, by D.R.C. Prescott. (2001)
- No. 27 The 2001 international piping plover census in Alberta, by D.R.C. Prescott. (2001)
- No. 28 Prairie rattlesnake (*Crotalus viridis viridis*) monitoring in Alberta – preliminary investigations (2000), by S.L. Rose. (2001)
- No. 29 A survey of short-horned lizard (*Phrynosoma hernandesi hernandesi*) populations in Alberta, by J. James. (2001)
- No. 30 Red-sided garter snake (*Thamnophis sirtalis parietalis*) education and relocation project – final report, by L. Takats. (2002)
- No. 31 Alberta furbearer harvest data analysis, by K.G. Poole and G. Mowat. (2001)
- No. 32 Measuring wolverine distribution and abundance in Alberta, by G. Mowat. (2001)
- No. 33 Woodland caribou (*Rangifer tarandus caribou*) habitat classification in northeastern Alberta using remote sensing, by G.A. Sanchez-Azofeifa and R. Bechtel. (2001)
- No. 34 Peregrine falcon surveys and monitoring in the Parkland Region of Alberta, 2001, by R. Corrigan. (2002)
- No. 35 Protocol for monitoring long-toed salamander (*Ambystoma macrodactylum*) populations in Alberta, by T. Pretzlaw, M. Huynh, L. Takats and L. Wilkinson. (2002)



- No. 36 Long-toed salamander (*Ambystoma macrodactylum*) monitoring study in Alberta: summary report 1998-2001, by M. Huynh, L. Takats and L. Wilkinson. (2002)
- No. 37 Mountain plover habitat and population surveys in Alberta, 2001, by C. Wershler and C. Wallis. (2002)
- No. 38 A census and recommendations for management for western blue flag (*Iris missouriensis*) in Alberta, by R. Ernst. (2002)
- No. 39 Columbian mountain amphibian surveys, 2001, by D. Paton. (2002)
- No. 40 Management and recovery strategies for the Lethbridge population of the prairie rattlesnake, by R. Ernst. (2002)
- No. 41 Western (*Aechmophorus occidentalis*) and eared (*Podiceps nigricollis*) grebes of central Alberta: inventory, survey techniques and management concerns, by S. Hanus, H. Wollis and L. Wilkinson. (2002)
- No. 42 Northern leopard frog reintroduction – year 3 (2001), by K. Kendell. (2002)
- No. 43 Survey protocol for the northern leopard frog, by K. Kendell. (2002)
- No. 44 Alberta inventory for the northern leopard frog (2000-2001), by K. Kendell. (2002)
- No. 45 Fish species at risk in the Milk and St. Mary drainages, by RL&L Environmental Services Ltd. (2002)
- No. 46 Survey of the loggerhead shrike in the southern aspen parkland region, 2000-2001, by H. Kiliaan and D.R.C. Prescott. (2002)
- No. 47 Survey of native grassland butterflies in the Peace parkland region of northwestern Alberta – 2001, by M. Hervieux. (2002)
- No. 48 Caribou range recovery in Alberta: 2001/02 pilot year, by T. Szkorupa. (2002)
- No. 49 Peace parkland native grassland stewardship program 2001/02, by A. Baker. (2002)
- No. 50 Carnivores and corridors in the Crowsnest Pass, by C. Chetkiewicz. (2002)
- No. 51 2001 Burrowing owl trend block survey and monitoring, Brooks and Hanna areas, by D. Scobie. (2002)
- No. 52 An evaluation of the ferruginous hawk population in Alberta based on recent trend data, by D.P. Stepnisky, G.L. Erickson, J. Iwaasa and B. Taylor. (2002)
- No. 53 Alberta amphibian call surveys. A pilot year. Final report, by L. Takats and C. Priestley. (2002)
- No. 54 Utilization of a roadside survey technique to survey burrowing owl (*Athene cunicularia hypugaea*) in southeastern Alberta, by J. Nicholson and C. Skiftun. (2002)

- No. 55 Alberta species at risk program and projects 2001-2002, by Alberta Sustainable Resource Development, Fish and Wildlife Division. (2002)
- No. 56 Developing a habitat-based population viability model for greater sage-grouse in southeastern Alberta, by C.L. Aldridge. (2001)
- No. 57 Peregrine falcon surveys and monitoring in the Northeast Boreal Region of Alberta, 2001, by R. Corrigan. (2002)
- No. 58 2002 burrowing owl trend block survey and monitoring, Brooks area, by R.F. Russell. (2002)
- No. 59 Rare plant inventory of the eastern edge of the lower foothills natural subregion, west-central Alberta, by J. Doubt. (2002)
- No. 60 Western (*Aechmophorus occidentalis*) and eared (*Podiceps nigricollis*) grebes of central Alberta: 2002 field summary, by S. Hanus, L. Wilkinson and H. Wollis. (2002)
- No. 61 Inventory of western spiderwort (*Tradescantia occidentalis*) in Alberta: 2002, by S. Peters. (2003)
- No. 62 Bullsnares (*Pituophis catenifer sayi*) in Alberta: literature review and data compilation, by K.J. Kissner and J. Nicholson. (2003)
- No. 63 Distribution of Ord's kangaroo rats in southeastern Alberta, by D.L. Gummer and S.E. Robertson. (2003)
- No. 64 Lethbridge prairie rattlesnake conservation project: 2002/2003 progress report, by R.D. Ernst. (2003)
- No. 65 Short-horned lizard (*Phrynosoma hernandesi hernandesi*) populations in Alberta – 2002 survey results, by J.D. James. (2003)
- No. 66 Inventory and monitoring protocol for naturally occurring western blue flag (*Iris missouriensis*) in Alberta, by R.D. Ernst. (2003)
- No. 67 The use of call playbacks for censusing loggerhead shrikes in southern Alberta, by D.R.C. Prescott. (2003)
- No. 68 Survey of bats in northeastern Alberta, by A. Hubbs and T. Schowalter. (2003)
- No. 69 Survey protocol for the Richardson's ground squirrel, by B.A. Downey. (2003)
- No. 70 Population estimates and a survey protocol for ferruginous hawks in Alberta, by B.N. Taylor. (2003)
- No. 71 Testing methods for detecting wolverine, by G. Mowat, C. Kyle and D. Paetkau. (2003)

- No. 72 A multi-species conservation strategy for species at risk in the Milk River basin: year 1 progress report, by R.W. Quinlan, B.A. Downey, B.N. Taylor, P.F. Jones and T.B. Clayton. (2003)
- No. 73 Long-toed salamander (*Ambystoma macrodactylum*) conservation in the Alberta foothills: 2002 field summary report, by L. Wilkinson and S. Hanus. (2003)
- No. 74 Researching Amphibian Numbers in Alberta (RANA): 2002 provincial summary, by L. Wilkinson and S. Hanus. (2003)
- No. 75 Distribution and habitat associations of the long-toed salamander (*Ambystoma macrodactylum*) in the Oldman River drainage, by K. Pearson. (2003)
- No. 76 Prairie rattlesnake (*Crotalus viridis*) hibernacula: monitoring history in Alberta 1987-2002, by K.J. Kissner and J. Nicholson. (2003)
- No. 77 Alberta species at risk program and projects 2002-2003, by Alberta Sustainable Resource Development, Fish and Wildlife Division. (2003)
- No. 78 Northern leopard frog reintroduction: Year 4 (2002), by K. Kendell. (2003)
- No. 79 Magrath Northern Leopard Frog Reintroduction Project - Year 1 Progress Report, by K.A. Romanchuk. (2003)
- No. 80 Conservation Overview of Butterflies in the Southern Headwaters at Risk Project (SHARP) Area, by N. Kondla. (2004)
- No. 81 Lethbridge Rattlesnake Conservation Project: 2003 Progress Report, by R. Ernst. (2004)
- No. 82 Shortjaw Cisco Species at Risk Assessment 2001, by M. Steinhilber. (2004)







NATIONAL LIBRARY OF CANADA  
Bibliothèque nationale du Canada



3 3286 53026763 8